


ADULT LAYER

preventative health care plan



Baycox Avian

For treatment of coccidia. Should only be used in chicks or hens off lay. Dose by beak or into the water (0.28mL per kg bodyweight) across 2 consecutive days. Refresh in between. WHP meat 14 days.

Aviverm

Dewormer. Dose by beak (beware toxicity) or into the water (1mL/9kg total bodyweight into 300mL water per adult bird), as a SINGLE dose. WHP meat 7 days, eggs 6 days

Flubenol

Dewormer. Add 1 level teaspoon (5g) to 7kg of dry feed and mix in well. Refresh daily for 7 days. Double the dose for Tapeworm or Gapeworm. No WHP

Spot-on 'Chicken Brew'

This product is off-label and cannot be guaranteed. Spot on the skin under the wing or back of the neck. Default WHP meat 63 days, eggs 10 days

The above plan is preventative for low-medium risk flocks, but if heavily burdened, high stocking rate, OR SICK your hen(s) will need to see a vet.

Recently rehomed ex-battery hens may require more frequent treatment, as they will have developed no immunity against worms and coccidia.

If you have few hens, adults and roaming a large area, you may not need to deworm at all. Ideally you would hand in fresh faecal samples to test, so we can tell you if they need to be dewormed, as unnecessary treatment can reduce how well our drugs will work in the future.

The above plan is for ADULT (>1yo) backyard flocks. Treat young chickens once every 3 months until one year old with Aviverm or Flubenol. Ensure chicks are on chick starter until at least 6 weeks old and are on chick starter for

chickens once every 3 months until one year old with Aviverm or Flubenol. Ensure chicks are on chick starter until at least 6 weeks old and are on chick starter for a time outside while they build immunity.



*Updated August 2025